

HIP ARTHROSCOPY

Post-Op Instructions

 www.TheAmericanOrthopedics.com/resources



Dr. Drew Ratner

DRESSING CHANGE

- Leave the dressing in place for 48 hours following surgery
- You may now apply a new, clean, dry dressing at least once a day.
 - Place sterile gauze over the incisions and then secure with tape
- After 5 days, you can leave the incisions open to air as long as they are dry and not draining.
- It is extremely important to keep the surgical site clean and dry at all times to help prevent infection.
- Signs of infection include redness, swelling, increased pain, warmth, fever (above 101.5 degrees F) and drainage from the wound. If you note any of these signs, please call the office immediately
- It is extremely important to keep the surgical site clean and dry at all times to help prevent infection.

BATHING

- You may shower **5 days after your surgery** and get your incisions wet as long as the wounds are dry and not draining. Let the water run down over the incisions but do not rub or scrub them.
- If the wounds are draining, then wait until they are dry before getting them wet.
- **Do not soak or submerge** the surgical site until the wounds are completely healed (typically 3-4 weeks after surgery).

ACTIVITY

- Rest and elevate your leg for the first 24 hours.
- Use a pillow under your calf and ankle. **Do not place a pillow under your knee or heel.**
- Use your crutches while walking. You may put your operative foot down for balance but are **not allowed to put more 50% of your weight on it.** You will typically be on crutches for 3 weeks post-operatively
- Do not let your **leg externally rotate for 3 weeks** post-operatively to protect your capsular repair
- **You will be discharged with a physical therapy prescription and a physical therapy protocol.** The protocol is a description for you and your physical therapist that details all of your restrictions and activities through your entire recovery. Please be sure to give this to your physical therapist.
- Please arrange your post-operative physical therapy appointment. Our office can help you with this if needed. You may start as soon as the day following your surgery. They will guide you in exercises to perform while in the clinic and at home. Avoid long periods of sitting without the leg elevated, or long-distance travel for 2 weeks

Call 864-807-9998 with any questions
about your post-op instructions.



PHYSICAL THERAPY

- You will be discharged with a physical therapy prescription and a physical therapy protocol. The protocol is a description for you and your physical therapist that details all of your restrictions and activities through your entire recovery. Please be sure to give this to your physical therapist.
- Please arrange your post-operative physical therapy appointment. Our office can help you with this if needed. You may start as soon as the day following your surgery, but most prefer the week after surgery. They will guide you in exercises to perform while in the clinic and at home.

STATIONARY BIKE

- If you have access to a stationary bike, it is good for your hip mobility to do 30 mins of stationary bike per day with no resistance and the seat lowered down such that you never fully extend the hip.

ICE THERAPY

- Icing is very important to decrease swelling and pain and improve mobility.
- Please use ice for 20 minutes each hour over surgical site.
- After 24 hours, continue to use the ice 3-4 times a day, 15-20 minutes each time.
- Be sure to keep clothing or a towel between ice and skin, as placing the ice directly on skin may cause frostbite. **DO NOT PLACE DIRECTLY ON SKIN.**

MEDICATIONS

- **If placed, nerve blocks and pain medication injected into the wound will wear off from 8-24 hours after surgery.**
- You will be given a narcotic pain medication. This should be taken as needed every 4-6 hours for pain. Take with a little food.
- Narcotic pain medications **will not alleviate all of your discomfort**
- **Ice and elevation** will aide in additional pain control
- Take medications as prescribed
- **Do not mix narcotics** with alcohol
- Common side effects of narcotic pain medications include
 - Nausea- Take medication with food. If prescribed you may use anti-nausea medication. If nausea persists, call the office to consider having the medication changed.
 - Drowsiness
 - Constipation - We recommend purchasing an over-the-counter stool softener (such as colace) to take as directed while taking narcotics to prevent constipation. You will be given a prescription for this.
- **Do not drive a vehicle or operate heavy machinery while on narcotic pain medications**
- Narcotic medications cannot be called into the pharmacy. They will require a hand written prescription from the office. Please monitor the amount of pain medications you have remaining in order to not run out.



HIP ARTHROSCOPY

Post-Op Instructions – American Orthopedics

864-807-9998

- Narcotics and pain medications will not be refilled on weekends or after hours.
- You will likely be given a prescription for aspirin 325mg to take post-operatively. This is taken only once a day for 4 weeks post-operatively to prevent blood clots. The dosing and duration may change on an individual basis, but it should be reflected in the prescription that you are given.
- You will also be given a prescription for **NAPROXEN 500mg** to be taken twice a day. This is really prescribed **to prevent extra bone from forming in your soft tissues (heterotopic ossification)** but also works to help minimize pain. Do not take if you are taking blood thinners or if instructed not to take the anti-inflammatories by your doctor. Take as directed. Take with food. Stop taking if you get any heartburn symptoms of stomach upset.

POST-OP OFFICE APPOINTMENT

- If the appointment was not arranged prior to surgery, please contact the office
- Your first post-operative visit will be scheduled in the office prior to surgery.
- Depending on the procedure plan for follow-up visits at 6 weeks, 3 months, and 6 months after surgery.

EMERGENCIES OR CONCERNS

- Email Dr. Ratner at drewratner@theamericanorthopedics.com
- Or Contact us or go to the nearest emergency room if you have any of the following:
 - Pain worsens or does not go away after pain medication is taken.
 - Any abrupt change in the nature or severity of pain
 - Redness or swelling in your thigh or calf that will not decrease.
 - Redness around incisions
 - Continuous drainage or bleeding from the incision (occasional drainage is expected)
 - Difficulties moving your fingers or extreme pain when moving your fingers
 - Numbness or tingling that does not go away or is getting worse
 - Cold foot or color change in extremity (Please be aware that the skin prep in surgery may have an orange, brown, or blue tint)
 - Develop a fever greater than 101.5 degrees F.
 - Chest pain, shortness of breath, intractable nausea/vomiting, or anything else concerning
 - If you have an emergency that requires immediate attention, proceed to the nearest emergency room.

**DREW
RATNER, MD**
ORTHOPEDIC SURGEON



864-807-9998



drewratner@theamericanorthopedics.com



www.TheAmericanOrthopedics.com



10 Enterprise Blvd. Ste 209
Greenville, SC 29615

Call 864-807-9998 with any questions
about your post-op instructions.



**AMERICAN
ORTHOPEDICS**