

# SMALL ROTATOR CUFF REPAIR & BICEPS TENOTOMY

## Rehab Protocols

 [www.TheAmericanOrthopedics.com/resources](http://www.TheAmericanOrthopedics.com/resources)



Dr. Drew Ratner

## PHASE I MAXIMUM PROTECTION (WEEKS 0-4)

### Goals

- Reduce inflammation
- Decrease pain
- Postural education
- PROM as instructed

### Exercise Progression

- UltraSling x 4 weeks
- Ice and modalities to reduce pain and inflammation.
- Cervical ROM and basic deep neck flexor activation (chin tucks).
- Instruction on proper head neck and shoulder (HNS) alignment.
- Active hand and wrist range of motion.
- Active shoulder retraction.
- Minimal Motion x 4 weeks (make sure can see hand at all times)
- Encourage walks and low intensity cardiovascular exercise to promote healing.

### Manual Intervention

- Scar tissue mobilization when incisions are healed.
- Graded GH mobilizations.

## PHASE II PROGRESSIVE STRETCHING & ACTIVE MOTION (WKS 4-8)

### Goals

- Discontinue sling.
- Postural education.
- Focus on posterior chain strengthening.
- Begin Passive / Active Assisted ROM:
  - Flexion 120°+
  - 20-30° ER @ 0° abduction
  - 30-45° ER at 70-90° abduction

Call **864-807-9998** with any questions  
about rehab protocols.



# SMALL ROTATOR CUFF REPAIR & BICEPS TENOTOMY

## Rehab Protocols – American Orthopedics

864-807-9998

### Exercise Progression

- Progress to full range of motion flexion and external rotation as tolerated. Use a combination of wand, pulleys, wall walks or table slides to ensure compliance.
- Gradual introduction to internal rotation using shoulder extensions (stick off back).
- Serratus activation; Ceiling punch (weight of arm) many initially need assistance.
- Scapular strengthening – prone scapular series (rows and I's). Emphasize scapular strengthening under 90°.
- External rotation on side (no resistance).
- Gentle therapist directed CR, RS and perturbations to achieve ROM goals.
- Cervical ROM as needed to maintain full mobility.
- DNF and proper HNS alignment with all RC/SS exercises.
- Low to moderate cardiovascular work. May add elliptical but no running

### Manual Intervention

- STM – global shoulder and CT junction.
- Scar tissue mobilization.
- Graded GH mobilizations.
- ST mobilizations.
- Gentle CR/RS to gain ROM while respecting repaired tissue.

## PHASE III – ADVANCED STRENGTHENING & ENDURANCE (WKS 8-12)

### Goals

- 90% passive ROM, 80-90% AROM by 12 weeks. Larger tears and patients with poor tissue quality will progress more slowly.
- Normalize GH/ST arthrokinematics.
- Activate RC/SS with isometric and isotonic progression.
- Continue to emphasize posterior chain strengthening but introduce anterior shoulder loading.

### Exercise Progression

- Passive and active program pushing for full flexion and external rotation.
- Continue with stick off the back progressing to internal rotation with thumb up back and sleeper stretch.
- Add resistance to ceiling punch.
- Sub-maximal rotator cuff isometrics (no pain).
- Advance prone series to include T's.
- Add rows with weights or bands.
- Supine chest-flys providing both strength and active anterior shoulder stretch.
- Supine (adding weight as tolerated) progressing to standing PNF patterns.
- Seated active ER at 90/90.
- Biceps and triceps PRE.
- Scaption; normalize ST arthrokinematics.
- 10 weeks; add quadruped or counter weight shift. Therapist directed RS and perturbations in quadruped – bilateral progressing to unilateral-tri pod position.

### Manual Intervention

- STM and Joint mobilization to CT junction, GHJ and STJ as needed.
- CR/RS to gain ROM while respecting repaired tissue.
- Manual perturbations.
- PNF patterns.

Call **864-807-9998** with any questions  
about rehab protocols.



### PHASE IV – ADVANCED STRENGTHENING & ENDURANCE (WKS 12-24)

#### PRE/PSE (weeks 12-20)

- Full range of motion all planes – emphasize terminal stretching with cross arm, TUB, triceps, TV, sleeper and door/pec stretch.
- Begin strengthening at or above 90° with prone or standing Y's, D2 flexion pattern and 90/90 as scapular control and ROM permit. Patient health, physical condition and goals/objectives will determine if strengthening above 90° is appropriate.
- Add lat pulls to gym strengthening program; very gradual progression with pressing and overhead activity.
- Continue with closed chain quadruped perturbations; add open chain as strength permits.
- Progress closed kinetic chain program to include push-up progression beginning with counter, knee then – gradual progression to full as appropriate.
- Initiate plyometric and rebounder drills as appropriate

#### Return To Sport Program

- Continue to progress RC and scapular strengthening program as outlined.
- Advance gym strengthening program.
- RTS testing for interval programs (golf, tennis etc.).
- Follow-up examination with the physician (6 months) for release to full activity.

#### Manual Intervention

- STM and Joint mobilization to CT junction, GHJ and STJ as needed.
- CR/RS to gain ROM while respecting repaired tissue.
- Manual perturbations.
- PNF patterns.

**DREW  
RATNER, MD**  
ORTHOPEDIC SURGEON



864-807-9998



drewratner@theamericanorthopedics.com



www.TheAmericanOrthopedics.com



10 Enterprise Blvd. Ste 209  
Greenville, SC 29615

**Call 864-807-9998 with any questions  
about rehab protocols.**



**AMERICAN  
ORTHOPEDICS**