

ANKLE ORIF Post-Op Instructions

🌐 www.TheAmericanOrthopedics.com/resources



Dr. Drew Ratner

INITIAL INSTRUCTIONS

- Ice along with pain medicine are your keys to initial pain management.
- After you arrive home, rest the remainder of the day. You may be tired or nauseated from the anesthesia and/or pain medication.
- You will be in a boot after surgery. You should wear this at all times after surgery.

DRESSING CHANGE

- Leave the dressing in place until your follow up appointment.
- It is extremely important to keep the surgical site clean and dry at all times to help prevent infection.
- Signs of infection include redness, swelling, increased pain, warmth, fever (above 101.4 degrees F) and drainage from the wound. If you note any of these signs, please call the office immediately

BATHING

- You may shower **after your surgery** but keep your dressing dry and covered.
- After your stitches are removed, you may shower and clean the wound gently with soap and water **but Do not soak or submerge** the surgical site until the wounds are completely healed (typically 3-4 weeks after surgery).

ACTIVITY

- May return to **sedentary work ONLY** the day after surgery, if pain is tolerable
- **OK to use a computer.**
- You will be non-weight bearing for 6 weeks and sometimes 8 weeks depending on the specifics of your surgery
- You can begin to work on range of motion of ankle immediately if pain allows

PHYSICAL THERAPY

- **You will be given a physical therapy prescription and a physical therapy protocol usually at your first postoperative appointment.** The protocol is a description for you and your physical therapist that details all of your restrictions and activities throughout your entire recovery. Please be sure to give this to your physical therapist.

Call 864-807-9998 with any questions
about your post-op instructions.



ANKLE ORIF

Post-Op Instructions - American Orthopedics

864-807-9998

CAM WALKER BOOT

- You will be put in a CAM Walker boot after surgery
- Boot should be worn at all times (except for hygiene and exercises)
- You will usually be in the boot for at least 6 weeks post-operatively, depending upon your procedure.

ICE THERAPY

- Icing is very important to decrease swelling and pain and improve mobility.
- Please use ice for 20 minutes at least 3 times per day.
- Be sure to keep clothing or a towel between ice and skin, as placing the ice directly on skin may cause frostbite. **DO NOT PLACE DIRECTLY ON SKIN.**

MEDICATIONS

- **If placed, nerve blocks and pain medication injected into the wound will wear off from 8-24 hours after surgery.**
- Recommendation is to take scheduled Tylenol 1000mg three times per day and 200mg Celebrex two times per day.
- You will be given a narcotic pain medication. This should be taken for breakthrough pain every 8 hours for pain if necessary. Take with a little food.
- **Ice and elevation** will aide in additional pain control
- Take medications as prescribed
- **Do not mix narcotics** with alcohol
- Common side effects of narcotic pain medications include
 - Nausea- Take medication with food. If prescribed you may use anti-nausea medication. If nausea persists, call the office to consider having the medication changed.
 - Drowsiness
 - Constipation - We recommend purchasing an over-the-counter stool softener (such as colace) to take as directed while taking narcotics to prevent constipation. You will be given a prescription for this.
- **Do not drive a vehicle or operate heavy machinery while on narcotic pain medications**
- **Narcotics and pain medications will not be refilled on weekends or after hours.**
- You will likely be given a prescription for aspirin 81mg twice per day to be taken post-operatively. This is taken only once a day for 4 weeks post-operatively to prevent blood clots. The dosing and duration may change on an individual basis, but it should be reflected in the prescription that you are given.
- You will be given Ergocalciferol (Vitamin D2) 50,000 units weekly to take for 8 weeks & Calcium 500mg to be taken three times per day for 8 weeks. Both of these medicines aid in bone healing.

Call 864-807-9998 with any questions
about your post-op instructions.



AMERICAN
ORTHOPEDICS

ANKLE ORIF

Post-Op Instructions - American Orthopedics

864-807-9998

POST-OP OFFICE APPOINTMENT

- Your first post-operative visit should be 10-14 days after surgery
- Depending on the specifics of the procedure, plan for additional follow-up visits at 6 weeks, 3 months, and 6 months after surgery.

EMERGENCIES OR CONCERNs

- Email Dr. Ratner at drewratner@theamericanorthopedics.com
- Or contact the office or go to the nearest emergency room if you have any of the following:
 - Pain worsens or does not go away after pain medication is taken.
 - Any abrupt change in the nature or severity of pain
 - Redness or swelling in your thigh or calf that will not decrease.
 - Redness around incisions
 - Continuous drainage or bleeding from the incision (occasional drainage is expected)
 - Difficulties moving your fingers or extreme pain when moving your fingers
 - Numbness or tingling that does not go away or is getting worse
 - Cold foot or color change in extremity (Please be aware that the skin prep in surgery may have an orange, brown, or blue tint)
 - Develop a fever greater than 101.5 degrees F.
 - Chest pain, shortness of breath, intractable nausea/vomiting, or anything else concerning
 - If you have an emergency that requires immediate attention, proceed to the nearest emergency room.

DREW
RATNER, MD
ORTHOPEDIC SURGEON

-  864-807-9998
-  drewratner@theamericanorthopedics.com
-  www.TheAmericanOrthopedics.com
-  10 Enterprise Blvd. Ste 209
Greenville, SC 29615

Call 864-807-9998 with any questions
about your post-op instructions.



AMERICAN
ORTHOPEDICS